

Fundamental Stellar Astrophysics Revealed at Very High Angular Resolution

arXiv:0905.1353v1 [astro-ph.SR] 8 May 2009

Contact: Jason Aufdenberg
(386) 226-7123

Embry-Riddle Aeronautical University, Physical Sciences Department
aufded93@erau.edu

Co-authors:

Stephen Ridgway (National Optical Astronomy Observatory)

Russel White (Georgia State University)

Introduction

A detailed understanding of stellar structure and evolution is vital to all areas of astrophysics. In exoplanet studies the age and mass of a planet are known only as well as the age and mass of the hosting star, mass transfer in intermediate mass binary systems lead to type Ia Supernova that provide the strictest constraints on the rate of the universe's acceleration, and massive stars with low metallicity and rapid rotation are a favored progenitor for the most luminous events in the universe, long duration gamma ray bursts. Given this universal role, it is unfortunate that our understanding of stellar astrophysics is severely limited by poorly determined basic stellar properties - effective temperatures are in most cases still assigned by blunt spectral type classifications and luminosities are calculated based on poorly known distances. Moreover, second order effects such as rapid rotation and metallicity are ignored in general. Unless more sophisticated techniques are developed to properly determine fundamental stellar properties, advances in stellar astrophysics will stagnate and inhibit progress in all areas of astrophysics. Fortunately, over the next decade there are a number of observational initiatives that have the potential to transform stellar astrophysics to a high-precision science. Ultra-precise space-based photometry from CoRoT (2007+) and Kepler (2009+) will provide stellar seismology for the structure and mass determination of single stars. GAIA (2011+) will yield precise distances to nearly a billion stars, providing accurate luminosities. However, *the unprecedented data from these upcoming missions will only translate to useful calibrations of stellar models if they are performed in concert with high angular resolution measurements provided by ground-based optical and infrared interferometers.*

Very high angular resolution (reaching < 1 milliarcsecond) observational astronomy at optical and infrared wavelengths is still in its infancy. Only in the last decade have multi-element interferometers become fully operational. In this white paper we highlight some of the incredible achievements made with optical and infrared interferometry over the last decade, and use these to emphasize the potential for these facilities, with continued support, to transform stellar astrophysics into a high-precision science.

Accurate Stellar Masses

Arguably the most fundamental parameter for a star is its mass, as this sets the timescale for evolution and determines its ultimate fate. Unfortunately mass is a difficult property to measure directly, and is typically only available for stars in binary systems with observable orbital motion. Mass estimates for the generic single stars must rely on mass-temperature or mass-luminosity calibrations, or worse, the predictions of untested stellar evolutionary models. The high resolution capabilities of optical interferometry have the potential to greatly increase the number and types of stars for which we have dynamical mass estimates, and greatly improve overall mass estimates.

The recent discoveries of young stars in nearby moving groups such as the β Pictoris

Association and the TW Hydrae Association (e.g. Zuckerman and Song 2004) provide many new opportunities to determine dynamical masses of young binaries (e.g. Boden *et al.* 2005a, 2007; Schaefer *et al.* 2008). Mass estimates at this early age are especially important because of the poorly understood input physics (e.g. convection, opacities) of pre-main sequence stars. With precise distance estimates to be provided by GAIA, relative orbits will provide dynamical masses of stars in rarer evolutionary states, such as those transitioning to the giant phases (e.g. Boden *et al.* 2005b), traversing the Hertzsprung Gap (Boden *et al.* 2006), and high-mass main sequence and Wolf Rayet stars (North *et al.* 2007a; Kraus *et al.* 2007). Only just recently have the prospects of high contrast imaging via non-redundant aperture masking on large-diameter telescopes been realized. This work can provide dynamical masses for substellar objects (Ireland *et al.* 2008, e.g.), a mass range where evolutionary models are very poorly constrained due to the age/temperature/mass degeneracies. As very high resolution interferometric techniques continue to mature, becoming more adaptable and sensitive, they will become an essential tool for determining stellar masses for stars spanning the entire H-R diagram.

Asteroseismology and Interferometry

While binary stars will continue to be the dominant device to determine stellar masses, density measurements provided by powerful asteroseismology measurements combined with radius measurements by optical interferometry offer a method to estimate masses for single stars. The photometric oscillations observed by asteroseismology reveal interior stellar properties through their dependence on the density (sound speed) distribution within the star. Since such distributions are a function of both mass and age, both of these fundamental stellar parameters are probed. Reliable masses and ages from asteroseismology require tight constraint of global stellar parameters, most importantly the stellar radius. Stellar radii from interferometry accurate to 3% when coupled with asteroseismology yield single star masses accurate to 4% (Cunha *et al.* 2007). In the case of the nearby subgiant β Hydri, this has been done to a precision of 2.8% (North *et al.* 2007b). Other examples of connections between asteroseismology and interferometry include a check on the mass of a δ Scuti star in the Hyades (Armstrong *et al.* 2006) and a radius and age for asteroseismology target τ Ceti (G8 V) (Di Folco *et al.* 2004). The synergy between ground-based interferometry and both ground-based programs (HARPS, CORALIE, ELODIE, UVES, UCLES, SIAMOUS, SONG) and space-based (MOST, CoRoT, WIRE, Kepler, PLATO) high-precision photometric missions will play a vital role in determining fundamental stellar masses.

Precise Radii and Temperatures

With available mass estimates, astronomers can begin the next step in high precision stellar astrophysics - testing the temperature and size predictions of stellar evolutionary mod-

els as a function of age. Historically the only stars that astronomers could measure accurate radii and (relative) temperatures of were in eclipsing binary systems (Andersen 1991). Long-baseline optical/infrared interferometry has changed this dramatically by providing sufficient resolution to measure the angular size of many nearby stars. Angular measurements with errors well under 1% are now commonplace. The limb-darkening corrections, normally modeled, are now subject to direct interferometric verification.

In order to convert a stellar angular diameter into a physical diameter, the distance must be known. Accurate binary orbits can provide this in some cases, but a more general solution is increasingly possible, with Hipparcos, GAIA and SIM providing/promising 1% distances to 10, 500 or 2500 pc. Interferometric + Hipparcos measurements of single M-dwarfs show that they are 10-15% larger than currently predicted by models (Berger *et al.* 2008), with a suggestion that the discrepancy increases with elevated metallicity. Examples of ultra-precise radii and temperatures (see Figure 1) have been measured for coeval binary stars (Kervella *et al.* 2008), metal poor population II stars (e.g. Boyajian *et al.* 2008), giant stars in the Hyades (Boyajian *et al.* 2009), and the enigmatic λ Boo itself (Ciardi *et al.* 2007).

Regarding stars with variable radii, a great achievement during the last decade has been the interferometric measurement of the pulsating diameters of seven classical cepheids: ζ Gem (Lane *et al.* 2000), η Aql, W Sgr, β Dor, I Car (Kervella *et al.* 2004a), δ Cep (Mérand *et al.* 2005), and Y Oph (Mérand *et al.* 2007). The precision of the δ Cep (see Figure 2) and Y Oph measurements was such that the projection factor, a value normally predicted by models and needed to correct radial velocity measurements, could be observationally constrained. GAIA distances will extend detailed study to a broad range of Cepheids.

The T_{eff} of a star is one of the small number of fundamental parameters of a stellar model. In order to confront models with real stars, the corresponding description of the star must be known. It is defined in terms of the luminosity and the radius by

$$T_{\text{eff}} = \left[\frac{L}{4\pi R^2 \sigma} \right]^{1/4} = \left[\frac{4f_{\text{bol}}}{\theta_{\text{LD}} \sigma} \right]^{1/4}$$

where f_{bol} is the bolometric flux, θ_{LD} is the angular diameter corrected for limb darkening, and σ is the Stefan-Boltzmann constant.

For a classical star with a well-defined surface, this lets us determine the T_{eff} by measuring the angular diameter and the observed flux, and modulo an understanding of any extinction, gives a precise answer. The determination of T_{eff} is typically limited by the photometry (or, with good photometry, by the absolute calibration of the photometry). In fact, with accumulating interferometric stellar measurements, it is now possible to turn the question around and, from stellar photometry alone, predict the angular diameters and T_{eff} of common spectral types to 1-2% (Kervella *et al.* 2004b). In the case of stars with a poorly defined surface (e.g. due to extended atmosphere, accretion disk, mass loss shell), an imaging capability allows a more detailed confrontation of observed and modeled brightness distributions (e.g.




Fig. 1.— Evolutionary tracks in the H-R diagram for 61 Cyg A (**left**) and B (**right**). The labels indicate the age in Gyr relative to the ZAMS. The rectangular box represents the classical $L - T_{\text{eff}}$ error box, and the diagonal lines represent the radius and its uncertainty determined by a combination of interferometry and the Hipparcos parallax (Kervella *et al.* 2008)




Figure 2 Angular diameter measurements showing the 5.36 day pulsation of δ Cephei with a model fit (A) and residuals (B) (Mérand *et al.* 2005). The solid curve is predicted from radial velocity data, where only the amplitude of the curve is adjusted to determine the projection factor.

Perrin *et al.* 2004b; Wittkowski *et al.* 2008) In the case of very hot stars, T_{eff} is particularly difficult to determine owing to the inability to observe the turnover of the Planck function in the far ultraviolet. With a measured angular diameter, a single photometric measure can give the surface brightness, and thanks to the simplicity of the spectrum, a reliable T_{eff} , and with distance a reliable luminosity, see Figure 3.

Limb Darkening

While precise masses and radii obtained from interferometric observations test and constrain stellar structure and evolution models, very high resolution observations of stellar photospheres test stellar atmosphere models, models that are vital for the construction of synthetic stellar spectral energy distributions and high-resolution synthetic spectra. Measurement of the center-to-limb intensity variation of a stellar photosphere probes the temperature structure of that atmosphere. This was first done for the Sun just over 100 years ago and helped to constrain models for the transport of energy in the solar atmosphere. Such studies are now possible for other stars.

Until recently, with the exception of Sirius (A1 V) Hanbury Brown *et al.* (1974) (Hanbury Brown, R. *et al.* 1974, MNRAS, 167,475), interferometric limb-darkening measurements have been limited to a small number of stars cooler than the sun: α Cas (K0 III) and α Ari (K2 III) (Hajian *et al.* 1998); Arcturus (K1.5 III) (Quirrenbach *et al.* 1996); Betelgeuse (M1Iab) (Perrin *et al.* 2004a); γ Sagittae (M0 III), V416 Lac (M4 III), and BY Boo (M4.5 III) (Wittkowski *et al.* 2001); ψ Phoenicis (M4 III) (Wittkowski *et al.* 2004).

In the last five years as long baselines have come on-line, hotter stars have been measured: Altair (A7 V) (Ohishi *et al.* 2004), Vega (A0 V) (Aufdenberg *et al.* 2006; Peterson *et al.* 2006b), α Cyg (A2 Ia) and β Ori (B8 Ia) (Aufdenberg *et al.* 2008). The interferometric confirmation that Vega is a pole-on, rapidly rotating star (see Figure 4) has recently driven astronomers to search for a replacement star, or set of stars, for defining photometric systems (Rieke *et al.* 2008).

Limb Darkening and Convection

High precision multi-wavelength angular diameter measurements reveal the wavelength dependence of limb-darkening. Such measurements of Procyon (Aufdenberg *et al.* 2005) and α Cen B (Bigot *et al.* 2006) provide tests of 3-D convective transport models. The interferometric data indicate a temperature gradient shallower than provided by the standard 1-D mixing-length convection. Further limb-darkening measurements, in particular at high spectral resolution (up to $R \sim 30,000$), will allow for more stringent tests and constraints on the




Fig. 3.— **Left:** The original Hanbury Brown intensity interferometer diameters together with the Hipparcos parallaxes, compared to massive star evolutionary tracks. **Right:** Expected error bars with SIM parallaxes and optical amplitude interferometry angular diameters, compared to evolutionary tracks.




Figure 4 A visibility curve for Vega (A0 V) showing the first and second lobes. The first lobe yields the angular size, while the second lobe contains limb-darkening information. In the case of Vega, a pole-on rapid rotator, the data show both limb and gravity darkening (solid line) (Aufdenberg *et al.* 2006). Such data are vital for probing the temperature structures of stellar atmospheres.

state-of-the-art multi-dimensional stellar atmosphere codes, those responsible for the recent substantial revision in the Sun's oxygen abundance (Asplund *et al.* 2004). *Global convective instabilities are present either in the core or envelope of most stars. Nevertheless, the appropriate modelling of convection remains one of the most difficult tasks in the context of stellar astrophysics.* - Cunha *et al.* (2007). High-resolution studies of stellar surfaces will play an important role in testing improved models for convection as they develop.

Stellar Rotation

Rotating stars offer a powerful tool for insight into stellar interiors. The mass distribution, opacities, and processes such as differential rotation and convection are normally lost in the spherical uniformity of normal stars. In rapid rotators, these factors contribute to the stellar shape and temperature distribution, which may follow more or less closely the idealizations of Roche and von Zeipel, and these factors will be implicated differently for stars of differing mass. A range of rotation rates for similar stars effectively constitutes a series of experiments. Interferometric imaging shows the distorted stellar shape. A first measurement of rotational deformation in α Eri (Domiciano de Souza *et al.* 2003) has already stimulated 14 publications of follow-up or interpretation. More recent imaging shows also the distribution of brightness temperature across the disk, and the limb darkening (eg. Zhao *et al.* 2009, in preparation, see Figure 5). For rapid rotators, of which Be stars are the classic example, interferometry can map the ejected material, which may be found in disks (Tycner *et al.* 2006) and/or polar winds (Kervella *et al.* 2009). Imaging may also serve an important role in understanding the interaction of rotation and pulsation, as in for example δ Sct stars (Peterson *et al.* 2006a) and other non-radial oscillators (Jankov *et al.* 2001).

Recent observations have spurred theoreticians to replace the Roche formalism (which uses a potential where the mass is a point source) with a more physically realistic model including a self-consistent gravitational potential and differential rotation (Jackson *et al.* 2005; MacGregor *et al.* 2007). These models predict that rotation significantly reduces the luminosity of young stars, by reducing the central core temperature, resulting, for example, in a $1.2 M_{\odot}$ star with a luminosity of $1.0 L_{\odot}$. Such models make matters worse for resolving the faint young Sun paradox (e.g. Minton and Malhotra 2007), where the standard solar model predicts the young Sun too cool to support life on the early Earth 4.0 Gyr ago, in contradiction to the geologic and fossil records. Testing the latest models of rapidly rotating stars with interferometric imaging will have an impact across stellar astrophysics.

High Angular Resolution Imaging in the Next Decade

Interferometric techniques already cover a large parameter space. In the near-IR, angular resolutions reach <1 milliarcsec, with single-measurement precision as fine as 20 microarcsec.




Fig. 5.— Model-independent images of rapid rotator stars α Aql (**left:** Monnier *et al.* (2007) and α Cep (**middle:** Zhao et al. 2009, in preparation) showing the shape effects of centripetal forces, and brightness distributions determined by polar brightening and equatorial darkening, polar axis projected angle, and limb darkening. **Right:** A Roche-von Zeipel model for α Oph based on interferometric data (Zhao et al. 2009, in preparation.)

Wavelength coverage extends over most of the atmospheric windows from 0.48 to 12.5 μm . Spectral resolutions are available up to 30,000 in R and I, 12,000 in J-H-K, adequate for characterizing most molecular bands and some individual spectral lines. In the N band, spectral resolutions up to 200 are well matched to grain and molecular opacity structure. Instrumentation and facility developments already underway will augment these capabilities somewhat, improve limiting sensitivity, and so forth. But the most dramatic development will be in the quality of imaging achieved. Most interferometric measurements are still made with only one telescope pair. Beam combination with 3, 4 or 6 telescopes is just in its infancy, but will rapidly come to be the preferred observing mode for the kinds of science described above, owing to its far more rapid accumulation of image information. Interferometry has important complementary potential for space, as well, for missions such as SIM, SI (Stellar Imager), and SPIRIT (Space Infrared Interferometric Telescope). The coming decade will be a period of rich development for high resolution stellar science.

REFERENCES

- Andersen, J. (1991). Accurate masses and radii of normal stars. *A&A Rev.*, **3**, 91–126.
- Armstrong, J. T., Mozurkewich, D., Hajian, A. R., Johnston, K. J., Thessin, R. N., Peterson, D. M., Hummel, C. A., and Gilbreath, G. C. (2006). The Hyades Binary θ^2 Tauri: Confronting Evolutionary Models with Optical Interferometry. *AJ*, **131**, 2643–2651.
- Asplund, M., Grevesse, N., Sauval, A. J., Allende Prieto, C., and Kiselman, D. (2004). Line formation in solar granulation. IV. [O I], O I and OH lines and the photospheric O abundance. *A&A*, **417**, 751–768.

- Aufdenberg, J. P., Ludwig, H.-G., and Kervella, P. (2005). On the Limb Darkening, Spectral Energy Distribution, and Temperature Structure of Procyon. *ApJ*, **633**, 424.
- Aufdenberg, J. P., Mérand, A., Foresto, V. C. d., Absil, O., Di Folco, E., Kervella, P., Ridgway, S. T., Berger, D. H., Brummelaar, T. A. t., McAlister, H. A., Sturmann, J., Sturmann, L., and Turner, N. H. (2006). First Results from the CHARA Array. VII. Long-Baseline Interferometric Measurements of Vega Consistent with a Pole-On, Rapidly Rotating Star. *ApJ*, **645**, 664–675.
- Aufdenberg, J. P., Ludwig, H.-G., Kervella, P., Mérand, A., Ridgway, S. T., Coudé Du Foresto, V., Ten Brummelaar, T. A., Berger, D. H., Sturmann, J., and Turner, N. H. (2008). Limb Darkening: Getting Warmer. In A. Richichi, F. Delplancke, F. Paresce, and A. Chelli, editors, *The Power of Optical/IR Interferometry: Recent Scientific Results and 2nd Generation*, pages 71–82.
- Berger, D. H., ten Brummelaar, T. A., Gies, D. R., Henry, T. J., McAlister, H. A., Merand, A., Sturmann, J., Sturmann, L., Turner, N. H., Aufdenberg, J. P., and Ridgway, S. T. (2008). The Radius-Luminosity Relation from Near-Infrared Interferometry: New M Dwarf Sizes from the CHARA Array. In G. van Belle, editor, *14th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun*, volume 384 of *Astronomical Society of the Pacific Conference Series*, pages 226–+.
- Bigot, L., Kervella, P., Thévenin, F., and Ségransan, D. (2006). The limb darkening of α Centauri B. Matching 3D hydrodynamical models with interferometric measurements. *A&A*, **446**, 635–641.
- Boden, A. F., Sargent, A. I., Akeson, R. L., Carpenter, J. M., Torres, G., Latham, D. W., Soderblom, D. R., Nelan, E., Franz, O. G., and Wasserman, L. H. (2005a). Dynamical Masses for Low-Mass Pre-Main-Sequence Stars: A Preliminary Physical Orbit for HD 98800 B. *ApJ*, **635**, 442–451.
- Boden, A. F., Torres, G., and Hummel, C. A. (2005b). Testing Stellar Models with an Improved Physical Orbit for 12 Bootis. *ApJ*, **627**, 464–476.
- Boden, A. F., Torres, G., and Latham, D. W. (2006). A Physical Orbit for the High Proper Motion Binary HD 9939. *ApJ*, **644**, 1193–1201.
- Boden, A. F., Torres, G., Sargent, A. I., Akeson, R. L., Carpenter, J. M., Boboltz, D. A., Massi, M., Ghez, A. M., Latham, D. W., Johnston, K. J., Menten, K. M., and Ros, E. (2007). Dynamical Masses for Pre-Main-Sequence Stars: A Preliminary Physical Orbit for V773 Tau A. *ApJ*, **670**, 1214–1224.
- Boyajian, T. S., McAlister, H. A., Baines, E. K., Gies, D. R., Henry, T., Jao, W.-C., O’Brien, D., Raghavan, D., Touhami, Y., ten Brummelaar, T. A., Farrington, C., Goldfinger, P. J., Sturmann, L., Sturmann, J., Turner, N. H., and Ridgway, S. (2008). Angular

- Diameters of the G Subdwarf μ Cassiopeiae A and the K Dwarfs σ Draconis and HR 511 from Interferometric Measurements with the CHARA Array. *ApJ*, **683**, 424–432.
- Boyajian, T. S., McAlister, H. A., Cantrell, J. R., Gies, D. R., Brummelaar, T. A. t., Farrington, C., Goldfinger, P. J., Sturmann, L., Sturmann, J., Turner, N. H., and Ridgway, S. (2009). Angular Diameters of the Hyades Giants Measured with the CHARA Array. *ApJ*, **691**, 1243–1247.
- Ciardi, D. R., van Belle, G. T., Boden, A. F., ten Brummelaar, T., McAlister, H. A., Bagnuolo, Jr., W. G., Goldfinger, P. J., Sturmann, J., Sturmann, L., Turner, N., Berger, D. H., Thompson, R. R., and Ridgway, S. T. (2007). The Angular Diameter of λ Boötis. *ApJ*, **659**, 1623–1628.
- Cunha, M. S., Aerts, C., Christensen-Dalsgaard, J., Baglin, A., Bigot, L., Brown, T. M., Catala, C., Creevey, O. L., Domiciano de Souza, A., Eggenberger, P., Garcia, P. J. V., Grundahl, F., Kervella, P., Kurtz, D. W., Mathias, P., Miglio, A., Monteiro, M. J. P. F. G., Perrin, G., Pijpers, F. P., Pourbaix, D., Quirrenbach, A., Rousselet-Perraut, K., Teixeira, T. C., Thévenin, F., and Thompson, M. J. (2007). Asteroseismology and interferometry. *A&A Rev.*, **14**, 217–360.
- Di Folco, E., Thévenin, F., Kervella, P., Domiciano de Souza, A., Coudé du Foresto, V., Ségransan, D., and Morel, P. (2004). VLTI near-IR interferometric observations of Vega-like stars. Radius and age of α PsA, β Leo, β Pic, ϵ Eri and τ Cet. *A&A*, **426**, 601–617.
- Domiciano de Souza, A., Kervella, P., Jankov, S., Abe, L., Vakili, F., di Folco, E., and Paresce, F. (2003). The spinning-top Be star Achernar from VLTI-VINCI. *A&A*, **407**, L47–L50.
- Hajian, A. R., Armstrong, J. T., Hummel, C. A., Benson, J. A., Mozurkewich, D., Pauls, T. A., Hutter, D. J., Elias, II, N. M., Johnston, K. J., Rickard, L. J., and White, N. M. (1998). Direct Confirmation of Stellar Limb Darkening with the Navy Prototype Optical Interferometer. *ApJ*, **496**, 484–+.
- Hanbury Brown, R., Davis, J., Lake, R. J. W., and Thompson, R. J. (1974). The effects of limb darkening on measurements of angular size with an intensity interferometer. *MNRAS*, **167**, 475–484.
- Ireland, M. J., Kraus, A., Martinache, F., Lloyd, J. P., and Tuthill, P. G. (2008). Dynamical Mass of GJ 802B: A Brown Dwarf in a Triple System. *ApJ*, **678**, 463–471.
- Jackson, S., MacGregor, K. B., and Skumanich, A. (2005). On the Use of the Self-consistent-Field Method in the Construction of Models for Rapidly Rotating Main-Sequence Stars. *ApJS*, **156**, 245–264.

- Jankov, S., Vakili, F., Domiciano de Souza, Jr., A., and Janot-Pacheco, E. (2001). Interferometric-Doppler imaging of stellar surface structure. *A&A*, **377**, 721–734.
- Kervella, P., Nardetto, N., Bersier, D., Mourard, D., and Coudé du Foresto, V. (2004a). Cepheid distances from infrared long-baseline interferometry. I. VINCI/VLTI observations of seven Galactic Cepheids. *A&A*, **416**, 941–953.
- Kervella, P., Thévenin, F., Di Folco, E., and Ségransan, D. (2004b). The angular sizes of dwarf stars and subgiants. Surface brightness relations calibrated by interferometry. *A&A*, **426**, 297–307.
- Kervella, P., Mérand, A., Pichon, B., Thévenin, F., Heiter, U., Bigot, L., Ten Brummelaar, T. A., McAlister, H. A., Ridgway, S. T., Turner, N., Sturmann, J., Sturmann, L., Goldfinger, P. J., and Farrington, C. (2008). The radii of the nearby K5V and K7V stars 61 Cygni A & B. CHARA/FLUOR interferometry and CESAM2k modeling. *A&A*, **488**, 667–674.
- Kervella, P., Domiciano de Souza, A., Kanaan, S., Meilland, A., Spang, A., and Stee, P. (2009). The environment of the fast rotating star Achernar. II. Thermal infrared interferometry with VLTI/MIDI. *A&A*, **493**, L53–L56.
- Kraus, S., Balega, Y. Y., Berger, J.-P., Hofmann, K.-H., Millan-Gabet, R., Monnier, J. D., Ohnaka, K., Pedretti, E., Preibisch, T., Schertl, D., Schloerb, F. P., Traub, W. A., and Weigelt, G. (2007). Visual/infrared interferometry of Orion Trapezium stars: preliminary dynamical orbit and aperture synthesis imaging of the θ^1 Orionis C system. *A&A*, **466**, 649–659.
- Lane, B. F., Kuchner, M. J., Boden, A. F., Creech-Eakman, M., and Kulkarni, S. R. (2000). Direct detection of pulsations of the Cepheid star ζ Gem and an independent calibration of the period-luminosity relation. *Nature*, **407**, 485–487.
- MacGregor, K. B., Jackson, S., Skumanich, A., and Metcalfe, T. S. (2007). On the Structure and Properties of Differentially Rotating, Main-Sequence Stars in the 1-2 $M_{\odot}$ Range. *ApJ*, **663**, 560–572.
- Mérand, A., Kervella, P., Coudé Du Foresto, V., Ridgway, S. T., Aufdenberg, J. P., Ten Brummelaar, T. A., Berger, D. H., Sturmann, J., Sturmann, L., Turner, N. H., and McAlister, H. A. (2005). The projection factor of δ Cephei. A calibration of the Baade-Wesselink method using the CHARA Array. *A&A*, **438**, L9–L12.
- Mérand, A., Aufdenberg, J. P., Kervella, P., Foresto, V. C. d., ten Brummelaar, T. A., McAlister, H. A., Sturmann, L., Sturmann, J., and Turner, N. H. (2007). Extended Envelopes around Galactic Cepheids. III. Y Ophiuchi and α Persei from Near-Infrared Interferometry with CHARA/FLUOR. *ApJ*, **664**, 1093–1101.

- Minton, D. A. and Malhotra, R. (2007). Assessing the Massive Young Sun Hypothesis to Solve the Warm Young Earth Puzzle. *ApJ*, **660**, 1700–1706.
- Monnier, J. D., Zhao, M., Pedretti, E., Thureau, N., Ireland, M., Muirhead, P., Berger, J.-P., Millan-Gabet, R., Van Belle, G., ten Brummelaar, T., McAlister, H., Ridgway, S., Turner, N., Sturmann, L., Sturmann, J., and Berger, D. (2007). Imaging the Surface of Altair. *Science*, **317**, 342–.
- North, J. R., Tuthill, P. G., Tango, W. J., and Davis, J. (2007a). γ^2 Velorum: orbital solution and fundamental parameter determination with SUSI. *MNRAS*, **377**, 415–424.
- North, J. R., Davis, J., Bedding, T. R., Ireland, M. J., Jacob, A. P., O’Byrne, J., Owens, S. M., Robertson, J. G., Tango, W. J., and Tuthill, P. G. (2007b). The radius and mass of the subgiant star β Hyi from interferometry and asteroseismology. *MNRAS*, **380**, L80–L83.
- Ohishi, N., Nordgren, T. E., and Hutter, D. J. (2004). Asymmetric Surface Brightness Distribution of Altair Observed with the Navy Prototype Optical Interferometer. *ApJ*, **612**, 463–471.
- Perrin, G., Ridgway, S. T., Coudé du Foresto, V., Mennesson, B., Traub, W. A., and Lacasse, M. G. (2004a). Interferometric observations of the supergiant stars α Orionis and α Herculis with FLUOR at IOTA. *A&A*, **418**, 675–685.
- Perrin, G., Ridgway, S. T., Mennesson, B., Cotton, W. D., Woillez, J., Verhoelst, T., Schuller, P., Coudé du Foresto, V., Traub, W. A., Millan-Gabet, R., and Lacasse, M. G. (2004b). Unveiling Mira stars behind the molecules. Confirmation of the molecular layer model with narrow band near-infrared interferometry. *A&A*, **426**, 279–296.
- Peterson, D. M., Hummel, C. A., Pauls, T. A., Armstrong, J. T., Benson, J. A., Gilbreath, G. C., Hindsley, R. B., Hutter, D. J., Johnston, K. J., Mozurkewich, D., and Schmitt, H. (2006a). Resolving the Effects of Rotation in Altair with Long-Baseline Interferometry. *ApJ*, **636**, 1087–1097.
- Peterson, D. M., Hummel, C. A., Pauls, T. A., Armstrong, J. T., Benson, J. A., Gilbreath, G. C., Hindsley, R. B., Hutter, D. J., Johnston, K. J., Mozurkewich, D., and Schmitt, H. R. (2006b). Vega is a rapidly rotating star. *Nature*, **440**, 896–899.
- Quirrenbach, A., Mozurkewich, D., Buscher, D. F., Hummel, C. A., and Armstrong, J. T. (1996). Angular diameter and limb darkening of Arcturus. *A&A*, **312**, 160–166.
- Rieke, G. H., Blaylock, M., Decin, L., Engelbracht, C., Ogle, P., Avrett, E., Carpenter, J., Cutri, R. M., Armus, L., Gordon, K., Gray, R. O., Hinz, J., Su, K., and Willmer, C. N. A. (2008). Absolute Physical Calibration in the Infrared. *AJ*, **135**, 2245–2263.
- Schaefer, G. H., Simon, M., Prato, L., and Barman, T. (2008). Preliminary Orbit of the Young Binary Haro 1-14c. *AJ*, **135**, 1659–1668.

- Tycner, C., Gilbreath, G. C., Zavala, R. T., Armstrong, J. T., Benson, J. A., Hajian, A. R., Hutter, D. J., Jones, C. E., Pauls, T. A., and White, N. M. (2006). Constraining Disk Parameters of Be Stars using Narrowband H α Interferometry with the Navy Prototype Optical Interferometer. *AJ*, **131**, 2710–2721.
- Wittkowski, M., Hummel, C. A., Johnston, K. J., Mozurkewich, D., Hajian, A. R., and White, N. M. (2001). Direct multi-wavelength limb-darkening measurements of three late-type giants with the Navy Prototype Optical Interferometer. *A&A*, **377**, 981–993.
- Wittkowski, M., Aufdenberg, J. P., and Kervella, P. (2004). Tests of stellar model atmospheres by optical interferometry. VLTI/VINCI limb-darkening measurements of the M4 giant ψ Phe. *A&A*, **413**, 711–723.
- Wittkowski, M., Boboltz, D. A., Driebe, T., Le Bouquin, J.-B., Millour, F., Ohnaka, K., and Scholz, M. (2008). J, H, K spectro-interferometry of the Mira variable S Orionis. *A&A*, **479**, L21–L24.
- Zuckerman, B. and Song, I. (2004). Young Stars Near the Sun. *ARA&A*, **42**, 685–721.